

Addressing Childhood Obesity for Type 2 Diabetes Prevention: Challenges and Opportunities

Deborah A. Galuska, Janelle P. Gunn, Ann E. O'Connor, and Ruth Petersen

■ **IN BRIEF** Addressing the problem of childhood obesity is an important component of preventing type 2 diabetes. Although children and their families ultimately make decisions about diet, physical activity, and obesity management, many groups have a role in making these choices easier. They do this by providing families with tools and resources and by implementing policies and practices that support a healthy diet and physical activity in the places where children and their families spend their time. Diabetes educators are an important part of the solution.

Among youth, diabetes is relatively rare; ~193,000 Americans (0.24%) <20 years of age have the diagnosis of either type 1 or type 2 diabetes (1). However, the incidence of type 2 diabetes has increased in youth. One study reported an annual adjusted increased incidence of 4.8% between 2002 and 2012 (2). The earlier an individual develops type 2 diabetes, the higher their lifetime risk of diabetes complications such as heart disease, vascular disease, vision loss, and amputation will be (3).

Although a number of causes may explain the increase in type 2 diabetes among youth, the increase in childhood obesity likely plays a major role. Obesity affects the body's ability to use insulin, therefore causing abnormal blood glucose levels (4). In the United States, childhood obesity doubled from 10% in 1999–2000 to 19% in 2015–2016, with >18 million children currently having obesity (5,6). This parallels the increase of type 2 diabetes in youth (2).

Type 2 diabetes is largely preventable. Addressing the problem of childhood obesity and its related risk factors of unhealthy diet and physical

inactivity is part of the solution and will have health benefits beyond the prevention of type 2 diabetes. In addition to being at higher risk for type 2 diabetes, children with obesity also are more likely to have other risk factors for chronic disease, such as high blood pressure and abnormal lipid levels (7,8). They also experience more bullying, stigmatization, and lower self-esteem (9,10). In addition, children with obesity are more likely to have obesity as an adult (11), and adult obesity is associated with a higher risk for type 2 diabetes (12), heart disease (12), some cancers (13), and complications during pregnancy (14).

Because of the complexity of the problem of childhood obesity, expert groups such as the Institute of Medicine and the Centers for Disease Control and Prevention (CDC) have long recognized that the solution would need a multi-level, multi-pronged approach. This approach involves multiple groups, including individuals; families; health care organizations; public health organizations; community settings; federal, state, and local agencies; and the private sector (15–17). This article briefly describes the risk factors for child-

Division of Nutrition, Physical Activity, and Obesity, Centers for Disease Control and Prevention, Atlanta, GA

Corresponding author: Deborah A. Galuska, dbg6@cdc.gov

<https://doi.org/10.2337/ds18-0017>

©2018 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0> for details.

hood obesity and the roles different groups can play in addressing obesity prevention and management. It closes by specifically identifying how diabetes educators can help reduce childhood obesity.

Preventing Unhealthy Weight Gain

Weight gain is a normal part of healthy development in children. Obesity prevention focuses on limiting weight gain to only what is needed for normal growth and optimal development. Children and families, as well as the communities in which they live, all have roles in obesity prevention.

Obesity prevention begins with children and their families. Starting early is key because children who are overweight as they enter kindergarten are four times more likely than their peers with a healthy weight to have obesity by eighth grade (18). Prevention activities can begin with a mother before a child is born. Prenatal exposures such as excessive maternal weight gain during pregnancy, gestational diabetes, tobacco use during pregnancy, and higher pre-pregnancy weight status all have been associated with childhood obesity (19–21).

Once a child is born, breastfeeding may also help prevent obesity; longer breastfeeding has been associated with a lower likelihood of childhood obesity in some, although not all, studies (22). Even as the evidence builds about the potential link between breastfeeding and healthy weight, breastfeeding has additional benefits for the mother and child and should be encouraged as the best first source of nutrition (23).

Maintaining healthy dietary patterns throughout childhood is an important obesity prevention strategy. As complementary foods are added to an infant's diet, an opportunity for establishing healthy eating patterns arises. Preferences and distinctions for sweet and salty tastes and foods begin in infancy (24).

As children transition to a “family diet,” caregivers' and family members' actions continue to shape their eating preferences and habits.

Dietary patterns consistent with the 2015–2020 Dietary Guidelines for Americans (DGA) are linked to better weight management, as well as improved health in general (25). These patterns include higher amounts of fruits, vegetables, and whole grains; moderate amounts of lean proteins; and lower amounts of processed foods, including those with added sugars, refined grains, and sodium. Resources such as the U.S. Department of Agriculture (USDA) MyPlate nutrition plan can guide parents in how to help their children's diet align with these patterns (26).

Whether specific foods and beverages contribute differentially to excess weight gain is not clear. The consumption of sugar-sweetened beverages has been linked to weight gain in children and adults (27); one review found that, among 17 foods or beverages, sugar-sweetened beverages were most consistently associated with excess weight gain (28). A more recent review found that, in general, the positive association is found when studies are limited to children <12 or to children <5 years of age (29).

In addition to poor diet, physical inactivity is another risk factor for childhood obesity (30). The 2008 Physical Activity Guidelines for Americans indicate that children ≥ 6 years of age need at least 60 min/day of physical activity for good health (31). Parents can help children be active by providing time for structured and unstructured activity, providing positive feedback, and encouraging children to try different types of physical activities (31).

Shorter sleep duration is a third risk factor for obesity (32). The recommended daily amount of sleep for optimal health ranges from 12 to 16 hours per 24-hour period for infants aged 4–12 months to 8–10 hours for teens aged 13–18 years (33). Parents can facilitate quality sleep through

establishing regular bedtimes and limiting the use of televisions and other screens shortly before bedtime and in the bedroom (34).

Obesity prevention efforts need to expand beyond the family, however. Despite their best efforts, families face barriers as they try to instill healthy habits in their children. These barriers include lack of time (35), lack of information about topics such as how to prepare healthy family meals (35), limited access to affordable healthy foods (35), and a lack of safe and accessible places for their children to play and be physically active (35,36). Furthermore, as children grow older, they spend increasing amounts of time in places outside of the home. The quality of these places can also make it easier or harder for children to eat healthier and be active. Practices and policies in childcare facilities, schools, and other community venues can reduce the barriers for children and their families in achieving and maintaining a healthy weight for children and are described in more detail below.

Childcare facilities and schools are important settings for supporting healthy behaviors because many children spend extended periods of time in them. Supports for healthy behaviors in childcare facilities and schools can be incorporated into individual facilities or across facilities through statewide policies and regulations such as licensing, staff training requirements, or state curriculum standards (15–17,37–39). For childcare facilities, states can also add obesity-related standards to quality rating and improvement systems (QRIS) to give facilities higher ratings when they incorporate activities and policies that promote a health weight; approximately one-half of states have included such standards in their QRIS (40).

A number of strategies can be employed in individual facilities to bring about system-level changes for healthier eating and physical activity. Individual childcare facilities and

schools can help children be more active by scheduling regular times for physical activity (16,37,39). In childcare settings, this can include designated time for structured and unstructured play, whereas in schools, this can include physical education classes and recess. Schools can also offer opportunities for physical activity outside of physical education. For example, they can offer intramural and extramural sports for youth at all levels of ability (37). Schools and other community venues can implement shared use agreements to increase access for physical activity for all community members (e.g., opening gyms, playgrounds, and tracks) (17). Childcare facilities and schools can encourage healthy eating consistent with the DGAs by serving healthier foods and beverages such as fruits and vegetables, whole grains, and lower-fat dairy products and limiting access to less healthy foods containing high amounts of sugar, sodium, or saturated fat (17,37–39). Changes can be made in foods and beverages offered in cafeterias, vending machines, and at special events and can address both meals and snacks. School curriculums can include information about diet and activity (37). Tools such as the Health Education Curriculum Analysis Tool can help schools determine whether their curriculums are likely to promote healthy behaviors (41). Both childcare facilities and schools can avoid using food or physical activity as a reward or punishment (37). In addition, teachers and staff in both facilities can model positive healthy behaviors.

Outside of childcare facilities and schools, the community at large also has a role in obesity prevention. Communities can help children be more active by providing places for activity and by making it safer and easier for children and families to walk or bike to these and other destinations (17,42,43). Examples of activity-friendly places for children include parks, playgrounds, and recre-

ation centers. Activity-friendly routes include well-maintained sidewalks and roads that have enforced speed limits, pedestrian-friendly intersections, and lanes that allow cyclists to ride safely. Many community groups, including those involved in planning, transportation, health, law enforcement, and education, have a role to play in making these changes happen. They can provide expertise, financial resources, programming, and space or help with interpreting and enforcing current laws and regulations (43).

To encourage healthy diets, communities can improve access to healthier and affordable foods, especially for individuals at highest risk for obesity, type 2 diabetes, and other chronic conditions. For example, community decision-makers can include economic or zoning incentives for supermarkets and farmers' markets to locate in low-income communities (16,17) and ensure that affordable public transportation to these places is available (16). In addition, government facilities embedded in communities can adopt healthy food policies that support access to healthy foods and beverages in their cafeterias, vending machines, and other food service venues (17,44).

These interventions to change the food environment, by themselves, may not change diet or weight status and may need to be combined with efforts to promote the new or improved venues and get people to purchase healthier choices once they come to the venue. For example, one study found that an intervention that added a new supermarket to a Philadelphia, Pa., neighborhood did not lead to changes in fruit and vegetable consumption or obesity for residents (45). The authors suggested that one reason for the finding was that the store was underused.

Like families, however, communities also experience barriers related to implementing these changes; these include limited understanding of what is needed to make changes

(46,47), limited resources (46–48), and competing priorities (46).

A number of CDC programs have helped states and communities address these barriers by providing financial resources and technical assistance. For example, through CDC funding, the Ohio Department of Health partnered with the Ohio Child Care Resources & Referral Association and Children's Hunger Alliance to provide more than 1,600 childcare centers, preschools, and home-based childcare programs with educational resources, training, and technical assistance on healthy eating and physical activity best practices for young children.

In addition to CDC programs, other federal programs and policies can also play a role in ensuring that children have access to healthier foods and opportunities for physical activity. For example, the USDA's National School Lunch Program and Child and Adult Food Program link the receipt of federal funding to the provision of healthier foods (49,50) and the implementation of school wellness policies (49).

Broader federal and state level policies can also be synergistic with local work. For example, only half of U.S. high school students participate in physical education at least once per week (51). Effectively addressing this gap will likely take a combination of policies at the government level that require specific amounts of physical education and adequate resources and training at the school level to implement the policies (16,52).

Obesity Management

Because of the large number of children with obesity, the problem of childhood obesity cannot be addressed solely through prevention efforts. Children with obesity often need focused and intensive intervention that starts in the health care setting.

In 2017, the U.S. Preventive Services Task Force (USPSTF) updated and confirmed a previous

recommendation that clinicians screen all children ≥ 6 years of age for obesity and, for children with obesity, offer or refer them to family-based comprehensive counseling and behavioral treatment interventions (i.e., pediatric weight management programs). These programs should have at least 26 contact hours over 2–12 months (53). Pediatric weight management programs can help children achieve a healthier weight by addressing the skills and efficacy of the child and the family to improve dietary patterns, increase physical activity, and improve sleep.

Uptake of the USPSTF recommendation has been limited by a small number of evidence-based programs across the country and struggles to fund programs in the long term. For example, in a 2013 survey of children's hospitals, only two-thirds of respondents with weight management programs indicated that the program met the USPSTF recommendation, and more than half of programs were operating at a financial loss (54). Implementation of the USPSTF guideline is also limited by family barriers such as readiness to participate in a program and the ability to schedule and get to multiple clinical appointments (55). Adequate coverage for services is also a barrier (56).

Recently, a group of cross-sector experts and stakeholders in obesity management identified strategies that could be used more effectively to move evidence-based interventions into routine clinical care. These strategies included integrating obesity treatment into a chronic care model, using multidisciplinary teams to deliver obesity interventions, ensuring that members of these teams have adequate training and resources, and developing payment models that cover children needing services (56).

Several programs also have the potential to improve the uptake of the USPSTF recommendation. For example, the Healthy Weight Partnership offers a program called “Mind, Exercise, Nutrition, Do It”

(MEND) in various settings (e.g., clinics, schools, and community centers) in the United States. The YMCA adapted MEND and is rolling out a version of this program called “Healthy Weight in Your Child” in YMCA sites across the country. This 4-month, family-based weight-management program focuses on helping children aged 7–13 years reach a healthy weight by working with them to improve their diet and increase their physical activity.

The CDC's Childhood Obesity Research Demonstration (CORD) project is testing models to address the needs of low-income families. CORD focuses on how to increase obesity screening, counseling, and referral to quality weight management programs for these high-risk families. Lessons learned from CORD will accelerate the expansion of similar programs.

Role of Diabetes Educators

In addition to the multiple groups identified above, diabetes educators can also be a part of solving the problem of childhood obesity. In working with people with diabetes, diabetes educators are likely already seeing children with obesity and helping them and their families achieve and maintain healthy weights. For example, many diabetes educators already bring in the families of patients to help inform and empower them to support the patient in managing their condition. Diabetes educators provide counseling to families on how to improve nutrition and physical activity. Examples include teaching families how to shop for and cook healthy foods and how to incorporate physical activity into daily routines. Although this counseling may be directed specifically at diabetes management, it can also help with weight management.

Diabetes educators can also link patients to community resources. Families can be struggling with a variety of health-related social needs that make it difficult for them to eat healthier or be more active. For exam-

ple, they may not have enough money for healthier foods or adequate transportation to get to a grocery store. They also may not be aware of places in their community where stores accept food assistance vouchers or where they can be physically active.

Diabetes educators can assess barriers to improving nutrition or physical activity through targeted questions about families' specific struggles, systematically identify a list of community resources that might help address their needs and the needs of the overall patient population, and refer families to these resources. Examples might include the locations of nearby parks, farmers' markets, or food banks or information on how to apply for the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) or other food assistance. The types and availability of these resources vary by community. However, information about them may be found by contacting local government offices or nonprofit groups; examining local, state, and federal websites; or using apps that identify nearby amenities. Active follow-up and problem-solving by diabetes educators can help patients address barriers they might experience when attempting to access these community resources (57).

In addition to their interactions with individual patients or small groups of patients, diabetes educators can also support obesity prevention in their community. As trusted community members, diabetes educators can educate decision-makers about their role in supporting the health of community members. For example, educators can learn about the health needs of their community by becoming familiar with findings from community needs assessments or other sources of community data. They can then look for opportunities to participate in community planning efforts that address needs such as safe and accessible roads and sidewalks for walking and bicycling or a lack of

local stores that sell healthier foods. They can talk with leaders in schools, childcare facilities, or worksites about the evidence-based programs and policies that could be implemented in these facilities (15–17,37–39,42,43).

Diabetes educators can also help make their own workplaces models for other community workplaces. For example, they can work with their worksite health coordinators to integrate nutrition standards for foods served and sold into facility policies and practices (44). This will help workers and patients have access to healthier foods while at the facility.

Conclusion

Addressing the problem of childhood obesity is an important component of preventing type 2 diabetes. Many groups have a role in addressing the problem through working with individual children and their families to improve diet and increase physical activity or making it easier for everyone to implement these healthy behaviors in the places where they spend their time. Diabetes educators are an important part of the solution.

Disclaimer

The findings and conclusions of this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Duality of Interest

No potential conflicts of interest relevant to this article were reported.

Author Contributions

D.A.G. and J.P.G. reviewed the literature and wrote the manuscript. All four authors contributed to discussion and reviewed/edited the manuscript. D.A.G. is the guarantor of this work and, as such, takes responsibility for the accuracy of the information presented.

References

1. Centers for Disease Control and Prevention. *National Diabetes Statistics Report, 2017: Estimates of Diabetes and Its Burden in the United States*. Available from www.cdc.gov/diabetes/pdfs/data/statistics/national-diabetes-statistics-report.pdf. Accessed 24 February 2018
2. Mayer-Davis EJ, Lawrence JM, Dabelea D, et al. Incidence trends of type 1 and type

- 2 diabetes among youths, 2002–2012. *N Engl J Med* 2017;376:1419–1429
3. Zoungas S, Woodward M, Li Q, et al. Impact of age, age at diagnosis and duration of diabetes on the risk of macrovascular and microvascular complications and death in type 2 diabetes. *Diabetologia* 2014;57:2465–2474
4. Al-Goblan AS, Al-Alfi MA, Khan MZ. Mechanism linking diabetes mellitus and obesity. *Diabetes Metab Syndr Obes* 2014;7:587–591
5. Hales CM, Carroll MD, Fryar CD, Ogden CL. Prevalence of obesity among adults and youth: United States, 2015–2016. *NCHS Data Brief* 2017;288:1–8
6. Ogden CL, Carroll MD, Lawman HG, et al. Trends in obesity prevalence among children and adolescents in the United States, 1988–1994 through 2013–2014. *JAMA* 2016;315:2292–2299
7. Güngör NK. Overweight and obesity in children and adolescents. *J Clin Res Periatr Endocrinol* 2014;6:129–143
8. Reilly JJ, Methven E, McDowell ZC, et al. Health consequences of obesity. *Arch Dis Child* 2003;88:748–752
9. van Geel M, Vedder P, Tanilon J. Are overweight and obese youths more often bullied by their peers? A meta-analysis on the correlation between weight status and bullying. *Int J Obes* 2014;38:1263–1267
10. Sikorski C, Luppá M, Luck T, Riedel-Heller SG. Weight stigma “gets under the skin”: evidence for an adapted psychological mediation framework: a systematic review. *Obesity* 2015;23:266–276
11. Simmonds M, Lewellyn A, Owen CG, Woolacott N. Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obes Rev* 2016;17:95–107
12. National Institutes of Health. *Managing Overweight and Obesity in Adults: Systematic Evidence Review from the Obesity Expert Panel, 2013*. Rockville, Md., U.S. Department of Health and Human Services, National Institutes of Health, 2013
13. Lauby-Secretan B, Scoccianti C, Loomis D, et al.; International Agency for Research on Cancer Handbook Working Group. Body fatness and cancer: viewpoint of the IARC Working Group. *N Engl J Med* 2016;375:794–798
14. Marchi J, Berg M, Dencker A, Olander EK, Begley C. Risks associated with obesity in pregnancy, for the mother and baby: a systematic review of reviews. *Obes Rev* 2015;16:621–638
15. Institute of Medicine. *Preventing Childhood Obesity: Health in the Balance*. Koplan JP, Liverman CT, Kraak VA, Eds. Washington, D.C., National Academies Press, 2005
16. Institute of Medicine. *Accelerating Progress in Obesity Prevention: Solving the Weight of the Nation*. Washington, D.C., National Academies Press, 2012

17. Centers for Disease Control and Prevention. Recommended community strategies and measurements to prevent obesity in the United States MMWR: Recomm Rep 2009;58(RR07):1–26
18. Cunningham SA, Kramer MR, Narayan KM. Incidence of childhood obesity in the United States. *N Engl J Med* 2014;370:403–411
19. Lau EY, Liu J, Archer E, McDonald SM, Liu J. Maternal weight gain in pregnancy and risk of obesity among offspring: a systematic review. *J Obes* 2014;2014:524939
20. Mamum AA, Mannan M, Doi SA. Gestational weight gain in relation to offspring obesity over the life course: a systematic review and bias-adjusted meta-analysis. *Obes Rev* 2014;15:338–347
21. Woo Baidal JA, Locks LM, Cheng ER, Blake-Lamb TL, Perkins ME, Taveras EM. Risk factors for childhood obesity in the first 1,000 days: a systematic review. *Am J Prev Med* 2016;50:761–779
22. Woo JG, Martin LJ. Does breastfeeding protect against childhood obesity? Moving beyond observational evidence. *Curr Obes Rep* 2015;44:207–216
23. Section on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics* 2012;129:e827–e841
24. Beauchamp GK, Mennella JA. Early flavor learning and its impact on later feeding behavior. *J Pediatr Gastroenterol Nutr* 2009;48(Suppl. 1):S25–S30
25. U.S. Department of Health and Human Services, U.S. Department of Agriculture. *2015–2020 Dietary Guidelines for Americans*. 8th ed. Washington, D.C., U.S. Department of Health and Human Services, 2015. Available from health.gov/dietaryguidelines/2015/guidelines. Accessed 24 February 2018
26. U.S. Department of Agriculture. ChooseMyPlate kids. Available from www.choosemyplate.gov/children. Accessed 24 February 2018
27. Malik VS, Pan A, Willett WC, Hu FB. Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. *Am J Clin Nutr* 2013;98:1084–1102
28. Fardet A, Boirie Y. Associations between food and beverage groups and major diet-related chronic diseases: an exhaustive review of pooled/meta-analyses and systematic reviews. *Nutr Rev* 2014;72:741–762
29. Frantsve-Hawley J, Bader JD, Welsh JA, Wright JT. A systematic review of the association between consumption of sugar-containing beverages and excess weight gain among children under age 12. *J Public Health Dent* 2017;77(Suppl. 1):S43–S66
30. Physical Activity Guidelines Advisory Committee. *Physical Activity Guidelines Advisory Committee Report, 2008*.

- Washington, D.C., U.S. Department of Health and Human Services, 2008
31. U.S. Department of Health and Human Services. *2008 Physical Activity Guidelines for Americans*. Washington, D.C., U.S. Department of Health and Human Services, 2008. Available from health.gov/paguidelines/pdf/paguide.pdf. Accessed 24 February 2018
32. Ruan H, Xun P, Cai W, He K, Tang Q. Habitual sleep duration and risk of childhood obesity: systematic review and dose-response meta-analysis of prospective cohort studies. *Sci Rep* 2015;5:16160
33. Hirshkowitz M, Whiton K, Albert SM, et al. National Sleep Foundation's updated sleep time duration recommendations: final report. *Sleep Health* 2015;1:233–243
34. American Academy of Pediatricians. Healthy sleep habits: how many hours does your child need? Available from www.healthychildren.org/English/healthy-living/sleep/Pages/Healthy-Sleep-Habits-How-Many-Hours-Does-Your-Child-Need.aspx. Accessed 26 February 2018
35. Ling J, Robbins LB, Hines-Martin V. Perceived parental barriers to and strategies for supporting physical activity and healthy eating among Head Start children. *J Community Health* 2016;41:593–602
36. Hesketh KR, Lakshman R, van Sluijs EMF. Barriers and facilitators to young children's physical activity and sedentary behaviour: a systematic review and synthesis of qualitative literature. *Obes Rev* 2017;18:987–1017
37. Centers for Disease Control and Prevention. School health guidelines to promote healthy eating and physical activity. *MMWR Recomm Rep* 2011;60(RR-5):1–76
38. Institute of Medicine. *Early Childhood Obesity Prevention Policies*. Washington, D.C., National Academies Press, 2011
39. National Resource Center for Health and Safety in Child Care and Early Education. *Achieving a State of Healthy Weight: 2014 Update*. Aurora, Col., University of Colorado Denver, 2014
40. Geary NA, Dooyema CA, Reynolds MA. Supporting obesity prevention in statewide quality rating and improvement systems: a review of state standards. *Prev Chronic Dis* 2017;14:E129
41. Centers for Disease Control and Prevention. Health education curriculum analysis tool. Available from www.cdc.gov/healthyyouth/hecat/index.htm. Accessed 24 February 2018
42. Community Preventive Services Task Force. Built environment approaches combining transportation system interventions with land use and environmental design. Available from www.thecommunityguide.org/sites/default/files/assets/PA-Built-Environments.pdf. Accessed 14 February 2018
43. U.S. Department of Health and Human Services. Step It Up! *The Surgeon General's Call to Action to Promote Walking and Walkable Communities*. Washington, D.C., U.S. Department of Health and Human Services, Office of the Surgeon General, 2015. Available from www.surgeongeneral.gov/library/calls/walking-and-walkable-communities/call-to-action-walking-and-walkable-communities.pdf. Accessed 24 February 2018
44. Food Service Guidelines Federal Workgroup. *Food Service Guidelines for Federal Facilities*. Washington, D.C., U.S. Department of Health and Human Services, 2017. Available from www.cdc.gov/obesity/downloads/guidelines_for_federal_concessions_and_vending_operations.pdf. Accessed 24 February 2018
45. Cummins S, Flint E, Matthews SA. New neighborhood grocery store increased awareness of food access but did not alter dietary habits or obesity. *Health Aff (Millwood)* 2014;33:283–291
46. Schuler BR, Saksvig BI, Nduka J, et al. Barriers and enablers to the implementation of school wellness policies: an economic perspective. *Health Promot Pract* 2018;Health Promot Pract 2018;19:873–883
47. Tomayko EJ, Prince RJ, Hoiting J, Braun A, LaRowe TL, Adams AK. Evaluation of a multi-year policy-focused intervention to increase physical activity and related behaviors in lower-resourced early care and education settings: Active Early 2.0. *Prev Med Rep* 2017;8:93–100
48. Deehr RC, Shumann A. Active Seattle: achieving walkability in diverse neighborhoods. *Am J Prev Med* 2009;37(6 Suppl 2):S403–S411
49. U.S. Department of Agriculture Food and Nutrition Service. National school lunch program and school breakfast program: nutrition standards for all foods sold in school as required by the Healthy, Hunger-Free Kids Act of 2010. *Fed Regist* 2017;81:50132–50151
50. Food and Nutrition Service, U.S. Department of Agriculture. Child and adult care food program. Available from www.fns.usda.gov/cacfp/child-and-adult-care-food-program. Accessed 14 February 2018
51. Kann L, McManus T, Harris WA, et al. Youth risk behavior surveillance—United States, 2017. *MMWR Surveill Summ* 2018;67(No. SS-8):1–114
52. Hills AP, Dengel DR, Lubans DR. Supporting public health priorities: recommendations for physical education and physical activity promotion in schools. *Prog Cardiovasc Dis* 2015;57:368–374
53. U.S. Preventive Services Task Force; Grossman DC, Bibbins-Domingo K, Curry SJ, et al. Screening for obesity in children and adolescents: U.S. Preventive Services Task Force recommendation statement. *JAMA* 2017;317:2417–2426
54. Children's Hospital Association. 2013 survey findings of children's hospitals: obesity services. Available from www.childrenshospitals.org/-/media/Files/CHA/Main/Issues_and_Advocacy/Key_Issues/Child_Health/Obesity/2013_survey_findings_obesity_services_report_050114.pdf. Accessed 24 February 2018
55. Perez AJ, Kebbe M, Holt NL, et al. Parent recommendations to enhance enrollment in multidisciplinary clinical care for pediatric weight management. *J Pediatr* 2018;192:122–129
56. Wilfley DE, Staiano AE, Altman M, et al. Improving access and systems of care for evidence-based childhood obesity treatment: conference key findings and next steps. *Obesity* 2017;25:16–29
57. Health Leads. Social determinants of health: overcoming the greatest barriers to patient care. Available from healthleadsusa.org/wp-content/uploads/2018/01/SDOH-White-Paper_Overcoming-the-Greatest-Barriers-to-Patient-Care_HL-and-UW.pdf. Accessed 24 February 2018